

Accessory cell functions in mononuclear cell cultures from uremic patients

J. LADEFOGED and E. LANGHOFF

Medical Department P, Division of Nephrology, Rigshospitalet, Copenhagen, Denmark

Accessory cell functions in mononuclear cell cultures from uremic patients. Mononuclear cells (PBMC) were isolated from dialyzed patients and healthy control subjects. Lymphocyte responses to stimulation with optimal and suboptimal concentrations of lectin (PHA), or stimulation with T cell receptor antibody (Leu 4) were found decreased in the patient cultures. The separate and the combined effects of exogenous interleukin-1 (IL-1) and interleukin-2 (IL-2) were examined in PHA and Leu 4 stimulated cell cultures. Addition of IL-1 did not normalize the decreased proliferation response of the patient cultures. In contrast, addition of IL-2 alone clearly enhanced and almost normalized the response of patient cultures stimulated with suboptimal concentrations of PHA. The combined addition of IL-1 and IL-2 gave no evidence of an additive effect of IL-1 and IL-2. Cell cultures from uremic and normal HLA-identical relative were examined. Substitution of uremic adherent monocytes with normal adherent monocytes as accessory cells did not improve the uremic T cell responses to stimulation with PHA. Furthermore, uremic adherent cells did not suppress the normal T cell responses. These results suggest that uremic accessory cells support T cell activation and, in particular, do not suppress T cell responses. The effect of IL-2 in the present study as well as previous findings of decreased IL-2 production in patients cultures may indicate that uremia primarily influences the proliferation of T cells.

In uremia a decreased immune response has been found in vivo [1, 2] and in vitro [3–6]. In vitro studies have shown that stimulated mononuclear cell cultures from hemodialyzed patients produce less interleukin-2 (IL-2) than normal cell cultures, whereas the expression of IL-2 receptors is unchanged or increased [6]. Activation of T cells is initiated by accessory cells which produce interleukin-1 (IL-1) and present lectins or antigens in the context of class II MHC products. Monocytes/macrophages are the most important accessory cells [7, 8]. Thus, impaired accessory cell function in uremia could be a possible mechanism for the decreased proliferation of uremic lymphocyte cultures. Therefore, the present study was undertaken to examine accessory cell functions in uremic patients.

Methods

Subjects

Peripheral venous blood was obtained before start of hemodialysis from 12 patients (4 females, 8 males) and from 12 (5

females, 7 males) age-matched control subjects. All patients had been on hemodialysis for at least one year and all had received blood transfusion. Predialysis blood samples were also obtained from nine uremic patients (2 females, 7 males) on hemodialysis and their healthy HLA-DR identical and mixed lymphocyte culture (MLC) negative relatives, either siblings or parents prior to kidney donation.

Patients and control subjects had no signs or symptoms of malignant diseases, inflammatory diseases, or were receiving immunosuppressive therapy.

Mitogen and antigen stimulated proliferation

Peripheral blood mononuclear cells (PBMCs) were isolated from the blood samples as previously described [5, 6].

Triplicates of 5×10^4 PBMCs were cultured in microtiter plates (Nunc, Denmark) in a total volume of 170 μ l. Phytohemagglutinin (PHA-P, Difco, Detroit, Michigan, USA) was used as lectin in the cultures in optimal (40 μ g/ml culture) and suboptimal (2 μ g/ml culture) concentrations. Leu 4 (anti-CD3) was used for direct stimulation of lymphocyte cultures at concentrations with maximal stimulatory effect found by titration (Leu 4; Becton Dickinson, Pennsylvania, USA, 1:20, 20 μ l/well). Lectin and Leu 4 stimulated cultures were incubated 72 hours in a humidified atmosphere with 5% carbon dioxide at 37° before 20-hour pulsing of the cultures with 14 C-thymidine (14 C-thymidine, 20 nCi/well, Amersham, UK) and harvested.

Part of the cultures was in parallel experiment supplemented with 1.25, 2.5 and 5 U/ml culture of purified human interleukin-1 (interleukin-1; Koch Light Ltd., UK) or purified human interleukin-2 (interleukin-2, Electronucleonics Inc., Maryland, USA, 10 μ l/well, or approximately 100 U).

Furthermore, six cultures from uremic patients and six controls were supplemented with normal erythrocytes. The erythrocytes were washed three times with saline and added to the culture to a final hematocrit of 1.5% in the culture medium before the PHA stimulation.

Isolation of cells for coculture experiments

Isolated PBMCs from nine patients and nine HLA-identical relatives were used in coculture experiment. Monocytes were obtained by two hour adherence in RPMI 1640 (Gibco) with 10% fetal calf serum (FCS) to plastic culture dishes. The adherent monocytes were detached from plastic after incubation for 30 minutes on ice and rinsed off the plates by ice-cold RPMI-1640 with 10% FCS. The nonadherent cell fraction was

Received for publication October 12, 1988

and in revised form June 19, 1989

Accepted for publication August 4, 1989

© 1990 by the International Society of Nephrology




Fig. 1. Mean ¹⁴C-thymidine uptake of 5×10^4 peripheral blood mononuclear cells (PBMCs) stimulated with optimal concentrations of PHA (40 µg/ml). Symbols are (▨) uremic PBMC cultures ($N = 12$), (□) normal PBMC cultures ($N = 12$). Vertical lines indicate SEM.

further purified twice over nylon wool columns to obtain a T-cell enriched cell fraction depleted of monocytes/macrophages [9]. T-cell enriched normal and uremic cell isolates (5×10^4 cells/well) were then cultured in the presence of 1% and 5% (500 and 2500 cells) normal or uremic adherent cells as accessory cells. As control 2.5×10^4 normal and 2.5×10^4 uremic PBMCs were cocultured. PHA was used at optimal (40 µg/ml cell culture) and suboptimal (2 µg/ml cell culture) concentrations. The culture conditions were as described above.

Esterase staining of monocytes/macrophages

Esterase staining was done as described by Tucker, Pierre and Jordan 1977 [10]. In one tube six drops of 1 mg% basic fuchsin (T01-0529, Technicon Instruments Corporation, New York, USA) was added to six drops of 7% sodium nitrite (T01-0530) to form the unstable hexazonium pararosanilin. Eighteen drops of 7% sodium cacodylate (T01-0531; buffer and serum lipase inhibitor) were added followed by three drops of alfa-naphthyl butyrate (T01-0680-81) as substrate. In a second tube, two drops of cells and two drops of fetal calf serum were mixed and added. The cells were then fixed for 40 seconds (fixative: 478.5 g diethyleneglycol, 50 ml formaldehyde, 10.9 g sodium acetate, and 0.6 g eserine sulphate in 1 liter distilled water). The contents of the second tube was then added to the first tube and incubated for two to three hours at 37°C before microscopic counting of dark red cells.

Statistical analysis

Student's *t*-test was used for comparison between groups. *P* values <0.05 were considered significant.

Results

Patient PBMCs stimulated with optimal concentrations of PHA (40 µg/ml) responded significantly lower than the control cultures. IL-1, IL-2, or IL-1 together with IL-2 (Fig. 1) had no additional effect on the response of the control cultures and did not enhance the response of the patient PBMCs. Suboptimal PHA stimulation (2 µg/ml) of the patient PBMC cultures

induced lower responses than in the control cultures ($P < 0.05$; Fig. 2). Addition of IL-1 alone improved patient as well as control PBMC responses, but the effect did not reach significance. In contrast, addition of IL-2 clearly enhanced both control and patient culture proliferation ($P < 0.01$) and decreased the difference between the two responses. No additive effects of IL-2 and IL-1 were observed. Results of stimulation with Leu 4 alone (Fig. 3) were comparable with those of suboptimal PHA stimulation. IL-1 alone had no effect on patient and control cultures, whereas IL-2 improved the PBMC responses ($P < 0.05$) in both uremic and control cultures. The PHA-2 and Leu 4 responses of uremic cell cultures with exogenous IL-2 were not significantly different from the normal cell cultures without exogenous IL-2 ($P > 0.1$).

Figure 4 shows the results of addition of erythrocytes to the culture medium. Addition of erythrocytes to the cultures improved the mitogen responses of patient and control cultures by about 20%, but did not normalize the decreased responses of the uremic cell cultures. The percentage of esterase positive cells (macrophages) was similar in patient and control PBMC cultures ($11.6 \pm 1.1\%$ [SEM], and $13.3 \pm 1.5\%$, $N = 11$, $P > 0.05$). In the coculture experiments, the mixed lymphocyte culture (MLC) responses were 1% of those obtained by PHA-stimulation, confirming the HLA-D identity between the nine patients and the nine healthy donors [11]. Stimulation of the patient and control PBMC cultures with 40 or 2 µg/ml PHA showed slightly lower responses of the uremic cell cultures, but the difference did not reach significance in this small material. However, there was a significant difference in the responses to 2 µg/ml PHA in the T-cell enriched fractions when normal and uremic patient cultures were compared (Fig. 5). Five percent adherent cells from patients or controls equally well enhanced T cell proliferation at suboptimal PHA concentrations (2 µg/ml, Fig. 5) and the same pattern was seen with PHA 40, but significant only for normal T cells (Fig. 6). Furthermore, there was no evidence that uremic adherent cells suppressed normal T cell responses to PHA, and no suppressive effect of uremic




Fig. 2. Mean ¹⁴C-thymidine uptake of 5×10^4 peripheral blood mononuclear cells (PBMCs) stimulated with suboptimal concentrations of PHA (2 µg/ml). Symbols are: (▨) uremic PBMC cultures (N = 11); (□) normal PBMC cultures (N = 11). Vertical lines indicate SEM.




Fig. 3. Mean ¹⁴C-thymidine uptake of 5×10^4 peripheral blood mononuclear cells (PBMCs) stimulated with optimal concentrations of Leu 4 (20 µl, 1:20). Symbols are: (▨) uremic PBMC cultures (N = 12); (□) normal PBMC cultures (N = 12). Vertical lines indicate SEM.

bulk PBMCs could be demonstrated when cocultured with normal bulk PBMCs.

Discussion

Previous studies [4, 5] have shown that PBMC cultures isolated from uremic patients respond lower than normal cultures to lectin stimulation as measured by DNA synthesis and production of IL-2. Expression of receptors for IL-2 has, however, been found normal in PHA-stimulated lymphocyte cultures from uremic patients [6] and cytotoxic effector cell functions that do not require cell proliferation remain nearly normal in uremia [12]. Antigen responses have also been found low [3, 13]. These findings of both in vivo and in vitro decreased antigen responses [14] suggest that accessory cell functions which initialize activation of T cells could be impaired. One mechanism could be a decreased production of IL-1 by the

accessory cells. This could eventually result in a decreased proliferation of activated T cells and a lower production of IL-2 as previously found in uremic cell cultures [4-6]. In this study we take advantage of purified preparations of IL-1 and IL-2 which have recently become available. We have furthermore studied accessory cell functions in coculture experiments of HLA-identical cell combinations. However, the present results give little support for a monocyte/macrophage defect. First, IL-1 did not reconstitute the patient cell responses whereas addition of exogenous IL-2 significantly improved the uremic cell responses. Second, coculture experiment showed that uremic macrophages/monocytes have the same capacity to serve as accessory cells for suboptimally stimulated T cells (Fig. 5). In these experiments a MLC stimulatory effect could be excluded since HLA-identical MLC negative combinations were used. Third, the percentage of esterase positive cells in the




Fig. 4. Effect of ^{14}C -thymidine uptake with or without addition of erythrocytes to uremic and normal peripheral lymphocytes cultures stimulated with PHA (40 $\mu\text{g}/\text{ml}$). Symbols are: (▨) uremia; (□) control. The difference between uremic and normals is significant ($P < 0.001$).




Fig. 5. Mean PHA (2 $\mu\text{g}/\text{ml}$) responses (^{14}C -thymidine uptake) in cocultures of isolated T cells from uremic patients with macrophages from their healthy HLA-D identical relatives, and vice versa. Hatched columns denote T cells alone or with 1% or 5% uremic macrophages (M), open columns are T cells supplied with normal macrophages. * $P < 0.01$ vs. uremic.

patient and the control PBMC cultures were similar and in accordance with the percentage published with this technique by Tucker, Pierre and Jordan [10].

Decreased in vitro transformation of uremic lymphocytes has in some studies been attributed to an enhanced suppressor activity by adherent cells (monocytes/macrophages) which are the predominating prostaglandin secreting cells [15]. In humans the results have been conflicting [16, 17], but in uremic rat models an enhanced suppressor activity by adherent cells has been reported [18]. Prostaglandins produced by macrophages have been shown to suppress T cell proliferation [19] and IL-2 production [20], but in the present study macrophages from patients showed no suppressive effect on the normal T-cell




Fig. 6. Mean PHA (40 $\mu\text{g}/\text{ml}$) responses (^{14}C -thymidine uptake) in cocultures of isolated T cells from uremic patients with macrophages from their healthy HLA-D identical relatives, and vice versa. Symbols are identical to Figure 5. * $P < 0.05$.

responses, and previous studies [6] have shown that blocking of PGE_2 production by indomethacin does not normalize the uremic lymphocyte responses. It cannot, however, be excluded that a suppressive effect of macrophages may be more pronounced in other lectin systems.

We cannot rule out that functions of other accessory cells are impaired in uremia. Dendritic cells (DC) are known to be active accessory cells in mice and in humans [7, 21, 22]. DC are plastic non-adherent cells and are therefore excluded in the present study by the plastic adherence procedure of the monocytes/macrophages.

In rare conditions impaired in vitro lymphocyte proliferation is caused by a deficiency of the enzyme system needed for the anaerobic glycolysis. This can be reversed by addition of an exogenous oxygen source [23, 24]. Addition of normal erythrocytes as oxygen suppliers improved the mitogen response of uremic and normal lymphocytes equally well, which show that the anaerobic condition that prevails in most culture medias does not especially injure uremic cells.

This and previous studies [5, 12] of immune responses in uremia confirm that the proliferative capacity of uremic T cells is impaired. Accessory cell functions as well as cytotoxic effector cell functions, however, seem normal.

Acknowledgment

This study was supported by the Danish Medical Research Council.

Reprint requests to J. Ladefoged, M.D., Medical Department P, Division of Nephrology, Rigshospitalet, 9 Blegdamsvej, 2100 Copenhagen, Denmark.

References

- KIRKPATRICK CH, WILSON WEC, TALMAGE DW: Immunologic studies in human organ transplantation: I. Observation and characterization of suppressed cutaneous reactivity in uremia. *J Exp Med* 119:727-742, 1964
- SELROOS O, PASTERNAK A, VIROLAINEN M: Skin test sensitivity and antigen-induced lymphocyte transformation in uremia. *Clin Exp Immunol* 14:365-370, 1973
- KUNORI T, FEHRMAN I, RINGDEN O, MOLLER E: In vitro characterization of immunological responsiveness of uremic patients. *Nephron* 26:234-239, 1980

4. KURZ P, KOHLRT H, MEUER S, HUTTEROTH T, ZUM BUSCHENFELDE K-HM: Impaired cellular immune responses in chronic renal failure: Evidence for a T cell defect. *Kidney Int* 29:1209-1214, 1986
5. LANGHOFF E, LADEFOGED J, ODUM N: Interleukin-2 (IL-2) enhancement of in vitro mitogen transformation of uremic lymphocytes. *Int Arch Allergy Appl Immunol* 81:5-11, 1986
6. LANGHOFF E, HOFMAN B, ODUM N, LADEFOGED J, PLATZ P, RYDER LP, SVEJGAARD A: Kinetic analysis of interleukin-2 (IL-2) production and expression of IL-2 receptors by uremic and normal lymphocytes. *Scand J Immunol* 25:29-36, 1987
7. INABA K, STEINMAN RM: Accessory cell-T lymphocyte interaction. Antigen-dependent and -independent clustering. *J Exp Med* 163:247-261, 1986
8. UNANUE ER, BELLER DI, LU CY, ALLEN PM: Antigen presentation: Comments on its regulation and mechanisms. *J Immunol* 132:1-5, 1984
9. WONG DM, MITTAL KK: HLA-DR typing: A comparison between nylon wool adherence and T cell rosetting in the isolation of B cells. *J Immunol Meth* 46:177-186, 1981
10. TUCKER SB, PIERRE RV, JORDAN RE: Rapid identification of monocytes in a mixed mononuclear cell preparation. *J Immunol Meth* 14:267-269, 1977
11. LANGHOFF E, JAKOBSEN BK, PLATZ P, RYDER LP, SVEJGAARD A: The impact of low donor specific MLR versus HLA-DR compatibility on kidney graft survival. *Transplantation* 39:18-21, 1985
12. LANGHOFF E, LADEFOGED J: In vitro natural killer and killer cell function in uremia. *Int Arch Allergy Appl Immunol* 78:218-220, 1985
13. CAPPEL R, BEERS DV, LIESNARD C, DRATWA M: Impaired humoral and cell-mediated immune response in dialyzed patients after influenza vaccination. *Nephron* 33:21-25, 1983
14. GOLDBLUM SE, REED WP: Host defenses and immunological alterations associated with chronic hemodialysis. *Ann Int Med* 93:597-613, 1980
15. KURLAND JI, BECKMANN R: Prostaglandin E production by human blood monocytes and mouse peritoneal macrophages. *J Exp Med* 147:941-957, 1978
16. OSAKI K, OTSUKA H, UOMIZU K, HARADA R, OTSUI Y, HASHIMOTO S: Monocyte-mediated suppression of mitogen responses of lymphocytes in uremic patients. *Nephron* 34:87-192, 1983
17. RUDDY MC, RUBIN AL, NOVOGRODSKY A, STENZEL KH: Decreased macrophage-mediated suppression of lymphocyte activation in chronic renal failure. *Am J Med* 75:571-579, 1983
18. RASKOVA J, RASKA K: Humoral inhibitors of the immune response in uremia. V. Induction of suppressor cells in vitro by uremic serum. *Am J Pathol* 111:149-155, 1983
19. CHOUAIB S, WELK K, MERTELSMANN R, DUPONT B: Prostaglandin E₂ acts at two distinct pathways of T lymphocyte activation: Inhibition of interleukin 2 production and down-regulation of transferrin receptor expression. *J Immunol* 135:1172-1179, 1985
20. RAPPAPORT RS, DODGE GR: Prostaglandin E inhibits the production of human interleukin 2. *J Exp Med* 155:943-948, 1982
21. YOUNG JW, STEINMAN RM: Accessory cell requirements for the mixed-leukocyte reaction and polyclonal mitogens, as studied with a new technique for enriching blood dendritic cells. *Cell Immunol* 111:167-195, 1988
22. LANGHOFF E, STEINMAN RM: Clonal expansion of human T lymphocytes initiated by dendritic cells. *J Exp Med* 169:315-320, 1989
23. PLATZ P, FABER V, HEDESKOV CJ, MORLING N, RYDER LP, SVEJGAARD A, THOMSEN M, THYME S, VELERIUS NH: Lymphocyte unresponsiveness to mitogens in vitro is not equivalent with severe immunodeficiency. *Birth Defects* 14:131-136, 1978
24. HEDESKOV CJ: Early effects of phytohaemagglutinin on glucose metabolism of normal human lymphocytes. *Biochem J* 110:373-380, 1968